Maximum CTDI 30

Note: All CT scans of the soft tissue neck should be performed with IV contrast. Except when the MD

specifically requests otherwise. If history is vocal cord paralysis, call referring MD for Chest

CT order.

PT Prep: Remove all metal objects from the neck to eliminate artifacts. Instruct the patient not to swallow

during the scan. Mark palpable mass(es) on patient's area of complaint when applicable.

Follow MCR IV contrast dosing guidelines.

Series 1: Scout AP and LAT-Supine "0" at Sternal Notch S 250 to I 100

Series 2: Helical scan of the neck from the anterior cranial fossa, about the level of the supra-orbital

rim to include pharynx, down to the top of the aortic arch.

Acquisition #1: Display scout and the prescribed slices from the anterior cranial fossa though the apices to the

top of the Aortic arch, 2.5m spacing and 2.5mm thickness.

Enhanced Scan: IV contrast at 2-3cc sec (follow MCR IV contrast dosing guidelines). Scan

delay of 45 sec.

Reconstruction: STANDARD AND BONE ALGORITHM with a DFOV large enough to include tip of the

nose and all of the mandible, approx 23-25cm.

Technique:

	750 HD 128 (room 2)	Optima 660 (32) OVIC	Optima 660 (room 1)	VCT 64 (room 3)
Noise Level	9.50	9.50	9.50	9.50
Interval	2.5mm	2.5mm	2.5mm	0.625
Axial/Helical Thickness	2.5mm	2.5mm	2.5mm	1.25
Pitch	1.375:1	1.375:1	1.375:1	1.375:1
Speed mm/rotation	55.00	27.50	55.00	55.00
Detector Rows			N/A	N/A
Detector Configuration			N/A	N/A
Beam Collimation	20mm	20mm	40mm	40mm
KV/mA	120KV/400mA	120KV/250mA	120KV/300mA	120KV/300mA
Scan Type	Helical Full 0.6 sec	Helical Full 1.0 sec	Helical Full 1.0sec	Helical 1.0 sec

Networking/ PACs: Send scouts

Send series 2 Standard Algorithm

Recon and send series 2 in a Bone Algorithm Recon and send lung images in Lung Algorithm

Reformat in Bone and Soft Tissue do sagittal and coronals and send to PACS.

Original Date: 4-8-04 Approved by: Dr. J. Campbell MCR

Revised Date: 11/9/04, 1/11/10, 9/16/10, 12/8/10, 01/28/13, 12/15/15, 08/18/20 Page 1 of 2

GE MDCT